

Date: November 2, 2009**Lesson Title:** 3.4 Linear Equations & the Intercept Form**Objective**

Write a linear equation in intercept form.

Explain the meaning of y-intercept and coefficient.

**IN**

When you analyze the walker graph, how do you know the starting value, the walkers direction and the rule (speed) of the walk?

Aug 26-9:45 AM

**Working Out with Equations
Investigation (25 minutes)**

Manisha starts her exercise routine by jogging to the gym. Her trainer says this activity burns 215 calories. Her workout at the gym is to pedal a stationary bike. This activity burns 3.8 calories per minute.

Oct 12-5:50 PM

First you'll model this scenario with your calculator.

Step 1:

Use calculator lists to write a recursive routine to find the total number of calories Manisha has burned after each minute she pedals the bike. Include the 215 calories she burned on her jog to the gym.

Manisha's Workout

Pedaling time (min) x	Total calories burned y
0	215
1	
2	
20	
30	
45	
60	

Oct 12-8:03 PM

Step 2:

Copy and complete the table using your recursive routine.

Step 3:

After 20 minutes of pedaling, how many calories has Manisha burned? How long did it take her to burn 443 total calories?



Manisha's Workout

Pedaling time (min) x	Total calories burned y
0	215
1	
2	
20	
30	
45	
60	

Oct 12-5:52 PM

Next you'll learn to write an equation that gives the same values as the calculator routines

Step 4:

Write an expression to find the total calories Manisha has burned after 20 minutes of pedaling. Check that your expression equals the value in the table.



Oct 17-2:33 PM

Step 5:

Write and evaluate an expression to find the total calories Manisha has burned after pedaling 38 minutes. What are the advantages of this expression over a recursive routine?



Oct 12-5:52 PM

Step 6:

Let x represent the pedaling time in minutes, and let y represent the total number of calories Manisha burns. Write an equation relating time to total calories burned.

Step 7:

Check that your equation produces the corresponding values in the table.

Oct 12-5:52 PM

Now you'll explore the connections between the linear equation and its graph.

Step 8:

Plot the points from your table on your calculator. Then enter your equation into the $Y =$ menu. Graph your equation to check that it passes through the points. Give two reasons why drawing a line through the points realistically models this

Oct 12-5:52 PM

The equation for Marisha's workout shows a linear relationship between the total calories burned and the number of minutes pedaling on the bike. You probably wrote this linear equation as

$$y = 215 + 3.8x \quad \text{or} \quad y = 3.8x + 215$$

The form $y = a + bx$ is the **intercept form**. The value of a is the **y-intercept**, which is the value of y when x is zero. The intercept gives the location where the graph crosses the y -axis. The number multiplied by x is b , which is called the **coefficient** of x .

Oct 17-8:41 PM

Example A:

Suppose Sam has already burned 325 calories before he begins to swim for his workout. His swim will burn 7.8 calories per minute.

- a. Create a table of values for the calories Sam will burn by swimming 60 minutes and the total calories he will burn after each minute of swimming.

Oct 17-2:35 PM

Example A solution:

- a. The total numbers of calories burned appear in the third column of the table. Each entry is 325 plus the corresponding entry in the middle column.

Sam's Swim

Swimming time (min)	Calories burned by swimming	Total calories burned
0	0	325
1	7.8	332.8
2	15.6	340.6
20	156	481
30	234	559
45	351	676
60	468	793



Oct 17-2:35 PM

Example A:

Suppose Sam has already burned 325 calories before he begins to swim for his workout. His swim will burn 7.8 calories per minute.

- b. Define variables and write an equation in intercept form to describe this relationship.
- c. On the same set of axes, graph the equation for total calories burned and the direct variation equation for calories burned by swimming.
- d. How are the graphs similar? How are they different?

Oct 17-2:35 PM

Example A solution:

- b. Let y represent the total number of calories burned, and let x represent the number of minutes Sam spends swimming.


$$y = 325 + 7.8x$$

- c. The direct variation equation is $y = 7.8x$. Enter it into Y1 on your calculator. Enter the equation $y = 325 + 7.8x$ into Y2. Check to see that these equations give the same values as the table by looking at the calculator table.

Oct 17-2:35 PM

Example A solution:

- d. The lower line shows the calories burned by swimming and is a direct variation. The upper line shows the total calories burned. It is 325 units above the first line because, at any particular time, Sam has burned 325 more calories. Both graphs have the same value of b , which is 7.8 calories per minute. The graphs are similar because both are lines with the same steepness. They are different because they have different y -intercepts.



What will different values of a in the equation $y = a + bx$ do to the graph?

Oct 17-2:35 PM

Topic Posters: (2 minutes)

What do you know about equations in the y-intercept form?

What do you have questions about?

On a Post-it-note:

Please write a response about how you feel about equations in the y-intercept form.

Aug 26-9:45 AM

Summary:

The equation $y = a + bx$ is called _____, since the value of

Out:

I have a test Tomorrow, so I will review 3.1-3.4

I will look on the Wiki under the Review section to find helpful study hints located there.

Aug 26-9:45 AM